

May 17, 2019

1010 2nd Ave South, PO Box 2806, Fargo, ND 58108
e-mail: kyle.aasand@ndsu.edu Phone: 241-5700

Kyle Aasand
Extension Agent/Cass County
Ag & Natural Resources

Cass County Palmer Amaranth Action Plan

A Guide for Landowners/Operators to Combat Palmer amaranth

Palmer amaranth is now on the North Dakota noxious weed list. If Palmer amaranth is suspected the following action plan will be taken by Cass County Weed Control and Cass County Extension and ask for the cooperation of landowners/operators.



Landowner/Operator

1. Begin scouting fields starting in late spring and continue through summer and fall, especially before harvest.
2. When a suspicious plant is found, contact the county extension agent or county weed officer.
 - Mark/flag the location and leave the plant in place to not spread seed. It is also easier to identify if it has not been pulled.
3. The Cass County Weed Officer and NDSU Extension agent will go to the field together and do the following:
 - Gather photos and information to send to NDSU weed specialist(s) for confirmation or ruling out Palmer amaranth.
 - Work with landowner/operator to develop a Palmer Amaranth Action Plan if Palmer amaranth is confirmed.
4. Landowner/operator should hand pull, bag on site and destroy confirmed Palmer amaranth plants.
5. Palmer amaranth plants should be burned or deeply buried to prevent movement of seed.
 - Palmer amaranth can germinate at depths up to 2 inches.
 - Burial depth should be below tillage level to prevent seeds from being brought up to the surface.
6. Work with the county weed officer and continue to survey the field at least 3-5 years after the weeds are removed. This helps to verify that no additional Palmer amaranth is found.
7. Please contact and cooperate with adjacent landowners/operators to help prevent the spread of Palmer amaranth to other land.

It is essential that all suspect plants be reported and identified; Palmer amaranth closely resembles other pigweed species and can be difficult to differentiate. One missed Palmer amaranth plant can produce enough seed to infest tens of acres. Landowners/operators that have an action plan in place and are working the plan will be in compliance with the noxious weed law. No citations will be issued if action plan is followed.

Contact Information:

Kyle Aasand

Cass County Extension

701-241-5700 (office) 701-650-8676 (cell)

Stan Wolf

Cass County Weed Officer

701-298-2388 (office) 701-730-6786 (cell)

North Dakota State University
Extension
PO Box 6050 Department 7070
Fargo, ND 58108-6050

CHANGE SERVICE REQUESTED


NonProfitOrg.U.S.
Postage

Paid

Permit No. 818
Fargo, N. Dak.

Identification of Palmer amaranth

Palmer amaranth looks very similar to both redroot pigweed and waterhemp. There are a few different characteristics that help distinguish these weeds. The simplest difference is the presence of short dense hairs on redroot pigweed as palmer and waterhemp have no hairs on the plants. The main distinguishing characteristic between palmer and waterhemp is the difference in petiole length. The petiole is the small stem that connects the leaf to the base stem of the plant and it is short on waterhemp, and on palmer it is as long or longer than the length of the leaf.



Wallet size Palmer amaranth ID cards are available by calling the Cass County Extension Office at 701-241-5700.



County commissions, North Dakota State University and U.S. Department of Agriculture cooperating. NDSU does not discriminate in its programs and activities on the basis of age, color, gender expression/identity, genetic information, marital status, national origin, participation in lawful off-campus activity, physical or mental disability, pregnancy, public assistance status, race, religion, sex, sexual orientation, spousal relationship to current employee, or veteran status, as applicable. Direct inquiries to Vice Provost, Title IX/ADA Coordinator, Old Main 201, 701-231-7708, ndsuoaa@ndsuo.edu.



Waterhemp and Palmer amaranth are two pigweed species that now are found in North Dakota and pose a serious threat to crop production. Both of these weeds have populations in other parts of the U.S. that have become resistant to nearly every herbicide mode of action used in crop production.

This publication will explain how the biology of waterhemp and Palmer amaranth, in addition to herbicide resistance, make them more difficult to control than redroot pigweed or Powell amaranth.

North Dakota farmers have dealt with pigweed (*Amaranthus*) for most of our farming history. Recent research has found that some populations of redroot pigweed are resistant to acetolactate synthase (ALS)-inhibiting (Group 2) herbicides in the state.

In the 1990s, waterhemp (*Amaranthus tuberculatus*) was identified in fields in the Red River Valley, and it has been rapidly spreading in the eastern half of North Dakota over the last decade.

Palmer amaranth (*Amaranthus palmeri*) was first identified in North Dakota in 2018. This weed has been one of the most difficult to control weeds in the southern United States for more than two decades. Populations found in North Dakota suggest that this plant can adapt and become competitive in the northern Great Plains as well.

Identification, Biology and Control of Palmer Amaranth and Waterhemp in North Dakota

Joe Ikley, NDSU Extension Weed Specialist

Brian Jenks, Weed Scientist, North Central Research Extension Center

Identification of Palmer amaranth and waterhemp

Proper identification is an important component of managing Palmer amaranth and waterhemp. Two common weeds that are mistaken for Palmer amaranth and waterhemp are redroot pigweed and Powell amaranth. Here are some tips to identify these four weeds from the seedling stage through plant maturity.

	Palmer amaranth (<i>Amaranthus palmeri</i>)	Waterhemp (<i>Amaranthus tuberculatus</i>)	Redroot pigweed (<i>Amaranthus retroflexus</i>)	Powell amaranth (<i>Amaranthus powellii</i>)
Stems	Smooth, no hair	Smooth, no hair	Short, dense hairs	Short, dense to sparse hairs
Petiole length	Long, as long or longer than leaf blade	Short	Typically short, can be long on mature plants	Typically short, can be long on mature plants
Inflorescence (Seed head)	Open, unbranched, very long Male are smooth, female are prickly	Open, unbranched Male and female are smooth	Compact, branched, 1-2" Smooth to rough, but not prickly	Compact, branched, 4-8" Smooth to rough, but not prickly
Bracts	Spiny bracts on female plants	No spiny bracts	No spiny bracts	No spiny bracts
Reproductive	Dioecious (separate male and female plants)	Dioecious (separate male and female plants)	Monoecious (male and female flowers on same plant)	Monoecious (male and female flowers on same plant)

Biology of Palmer amaranth and waterhemp

Palmer amaranth (Figure 1) and waterhemp (Figure 5) do not have hairs on the stems. Redroot pigweed (Figure 9) has short, dense hairs, while Powell amaranth (Figure 12) can be sparsely to densely hairy. Always look for hairs at the top of the plant near the newest growth.

Pull older, more mature leaves of suspected Palmer amaranth or waterhemp plants to examine petiole length. Palmer amaranth (Figure 4) has petioles as long as or longer than the leaf blade. Waterhemp (Figure 8) petioles are shorter than the leaf blades.

Palmer amaranth female plants are easily distinguished at maturity from other pigweeds. There are spiny bracts (Figure 15) at each leaf axil, and the seed head (Figure 16) is prickly and rough to handle. Male plants have smooth inflorescence that can be confused with other pigweeds.

Palmer amaranth and waterhemp have smooth stems at maturity. Palmer amaranth (Figure 2, Figure 3) has the longest inflorescence/seed heads of the pigweeds. The long terminal seed head often can reach 2 to 3 feet long by late summer.

Waterhemp (Figure 6, Figure 7) tends to have thin inflorescence/seed heads compared to the other pigweeds.

Redroot pigweed and Powell amaranth have hairy stems near the inflorescence/seed heads. The seed heads of redroot pigweed and Powell amaranth are compact and branched. Redroot pigweed (Figure 10, Figure 11) has short branches, while Powell amaranth (Figure 13, Figure 14) has longer branches.



■ Genetic Diversity

Both Palmer amaranth and waterhemp easily adapt to new locations. These weeds are both dioecious, meaning they have separate male and female plants. This means that plants must outcross to pollinate and produce seed, which leads to increased genetic diversity. Research has shown that herbicide-resistance traits can transfer on pollen, which quickly can expand geographic areas of resistant populations.

■ Prolific Seed Production

Both Palmer amaranth and waterhemp can produce up to 100,000 seeds per female plant in direct competition with crops. In the absence of competition, plants can produce more than 1 million seeds per plant. This prolific seed production allows for populations to quickly take over fields if plants are not controlled and are allowed to produce seed.

■ Small Seed Size

The seeds of all pigweed species are very small, which allows seed to be easily transported through various means. Palmer amaranth and waterhemp have been introduced to new areas through contaminated straw, hay and other animal feed. Equipment, especially combines, can carry seed long distances. Contamination of Conservation Reserve Program seed and cover crop seed have led to new introductions in the Corn Belt. Wildlife and migratory birds also can spread seed.

■ Extended Germination Period

Both Palmer amaranth and waterhemp can germinate throughout the growing season. Germination typically begins mid-May in North Dakota, but can continue throughout the summer into September. This leads to management challenges in crops that struggle to canopy and could be a challenge after harvesting early maturing crops.

■ Competitiveness

Palmer amaranth and waterhemp can be aggressive competitors under ideal growth conditions. These plants can grow 2 to 3 inches per day if conditions are favorable. Postemergence herbicides are most effective when the plants are smaller than 4 inches. Their rapid growth can lead to compressed spray application windows for optimal control. Weeds larger than 4 inches likely cannot be controlled by any post-emergent herbicide, including dicamba or 2,4-D.

■ Herbicide Resistance

Resistance to the ALS-inhibiting (Group 2) herbicides and glyphosate (Group 9) is so widespread in both plants that it is safe to assume resistance to both modes of action. Protoporphyrinogen oxidase (PPO)-inhibiting (Group 14) herbicide resistance is becoming more common. Populations also have been identified with resistance to dinitroanilines (Group 3), growth regulators (Group 4), triazines (Group 5), chloroacetamides (Group 15), and 4-Hydroxyphenylpyruvate dioxygenase (HPPD)-inhibitors (Group 27).



Figure 1. Palmer amaranth stem



Figure 5. Waterhemp stem



Figure 9. Redroot pigweed stem



Figure 12. Powell amaranth stem



Figure 2. Palmer amaranth seed head



Figure 6. Waterhemp seed head



Figure 10. Redroot pigweed seed head



Figure 13. Powell amaranth seed head



Figure 3. Palmer amaranth seed head



Figure 7. Waterhemp seed head



Figure 11. Redroot pigweed seed head



Figure 14. Powell amaranth seed head



Figure 4. Palmer amaranth petiole



Figure 8. Waterhemp petiole



Figure 15. Palmer amaranth spiny bracts



Figure 16. Palmer amaranth seed head

Management of Palmer amaranth and waterhemp

■ Crop Rotation

Palmer amaranth and waterhemp are very competitive against most summer annual crops, but the availability of effective herbicides for corn and soybeans provide more options for chemical weed control. Small grains offer a competitive advantage compared to row crops due to early row closure that helps prevent late-season germination. Limited post-emergent herbicides for pulse crops, flax, canola, sunflowers and other minor crops make Palmer amaranth and waterhemp control almost a no-win situation in these crops.

Effective crop and chemical rotation can help preserve effective herbicides and reduce the chances of Palmer amaranth and waterhemp developing resistance to herbicides. Due to constantly evolving herbicide products and the wide variety of crops across the state, specific herbicide programs are updated regularly at <https://www.ag.ndsu.edu/weeds>.

■ Overlapping Residual Herbicides

The extended germination period of Palmer amaranth and waterhemp make these two weeds more difficult to control than traditional pigweeds. In addition to the use of multiple effective modes of action, multiple applications of residual herbicides should be the foundation of a herbicide program focused on Palmer amaranth or waterhemp control.

Research has shown that effective herbicide programs include multiple applications of residual herbicides. Apply residual herbicides at planting to control early emerging weeds. An in-crop application of effective post emergence herbicide, plus an additional residual herbicide about 28 days after planting, can help reduce weed pressure until the crop can canopy.

For most crops, the most effective residual herbicides are the Group 15 herbicides, which help control the later-emerging plants.

■ Good Agronomic Practices

Season-long control of Palmer amaranth and waterhemp relies on the ability to get a crop canopy. This means that good agronomic practices are an important component of a management program. Problems such as soybean cyst nematode (SCN) and iron-deficiency chlorosis (IDC) should be addressed because these problems can delay or prevent the formation of a full crop canopy.

All photos by Joseph Ikley, Brian Jenks and Tom Peters, North Dakota State University

NDSU Extension does not endorse commercial products or companies even though reference may be made to tradenames, trademarks or service names. NDSU encourages you to use and share this content, but please do so under the conditions of our Creative Commons license. You may copy, distribute, transmit and adapt this work as long as you give full attribution, don't use the work for commercial purposes and share your resulting work similarly. For more information, visit www.ag.ndsu.edu/agcomm/creative-commons.

For more information on this and other topics, see www.ag.ndsu.edu/extension

County commissions, North Dakota State University and U.S. Department of Agriculture cooperating. NDSU does not discriminate in its programs and activities on the basis of age, color, gender expression/identity, genetic information, marital status, national origin, participation in lawful off-campus activity, physical or mental disability, pregnancy, public assistance status, race, religion, sex, sexual orientation, spousal relationship to current employee, or veteran status, as applicable. Direct inquiries to Vice Provost for Title IX/ADA Coordinator, Old Main 201, NDSU Main Campus, 701-231-7708, ndsu.eoaa@ndsu.edu. This publication will be made available in alternative formats for people with disabilities upon request, 701-231-7881. 1M-5-19

■ Tillage

Thorough tillage controls emerged plants prior to planting. Inter-row cultivation also can control emerged plants in-crop. Deep tillage can help control areas with a large population of emerged plants.

The small seed size of Palmer amaranth and waterhemp restricts germination to the top one-half inch of the soil profile. If seed is buried deeper than a few inches, Palmer and waterhemp cannot emerge. Deep tillage should be a one-time event to avoid bringing buried seed back near the surface. Pigweed seed can be expected to remain viable at least 3 to 5 years on the soil surface. Research has shown that buried seed can remain viable for up to 16 years.

■ Cereal Rye Cover Crop

Research in southern areas of the U.S. has shown promise with using a cereal rye cover crop to suppress Palmer amaranth and waterhemp. About 1,500 pounds of cereal rye biomass per acre at termination can delay the time it takes Palmer amaranth to grow to 4 inches, by about 2 weeks. Producing enough biomass for complete weed control in North Dakota is unlikely, but it can help widen the window to make a post emergence herbicide application.

■ Hand Weed

Hand weeding is time and labor intensive, but it can be very effective in a total weed management program. A zero-tolerance policy for seed production can help with management of Palmer amaranth and waterhemp due to their abundant seed production. If you suspect a pigweed is Palmer amaranth or waterhemp, pull the plants to help prevent a seed rain and avoid major issues in future years.

■ Field Edges, Borders, Ditches and Fence Rows

Palmer amaranth and waterhemp can live and produce seed in areas beyond crop fields. Field edges and borders between fields are important areas to monitor for weeds. Pigweeds in these areas can contribute to spreading resistant populations since herbicide resistance can spread via pollen.

Communication with neighbors helps prevent pollen shed and seed production in these areas. Weeds also can thrive in ditches and fencerows. Pasture and right-of-way herbicides can be effective against Palmer amaranth and waterhemp.

■ Sanitation

Equipment is one of the easiest ways to move pigweed seed from one area to another. If you have infested fields, farm these fields last. Combines are especially capable of spreading seed, so harvest infested fields last. If you run a combine through a patch of Palmer amaranth or waterhemp, you will spread the seed to other fields. Since the seed size is very small, thoroughly clean equipment to ensure seed is not spread to additional fields.